

Approximating the satisfiability threshold for random k -XOR-formulas

Nadia CREIGNOU*

Laboratoire d'Informatique de Marseille
LIM, FRE CNRS 2246
Université de la Méditerranée
Marseille, France

Hervé DAUDÉ †

Centre de Mathématiques et d'Informatique
LATP, UMR CNRS 6632
Université de Provence
Marseille, France

Olivier DUBOIS‡

CNRS-Université de Paris 6
LIP6
Paris, France

February 1, 2008

Abstract

In this paper we study random linear systems with k variables per equation over the finite field $GF(2)$, or equivalently k -XOR-CNF formulas. In a previous paper Creignou and Daudé proved that the phase transition for the consistency (satisfiability) of such systems (formulas) exhibits a sharp threshold. Here we prove that the phase transition occurs as the number of equations (clauses) is proportional to the number of variables. For any $k \geq 3$ we establish first estimates for the critical ratio. For $k = 3$ we get 0.93 as an upper bound, 0.89 as a lower bound, whereas experiments suggest that the critical ratio is approximately 0.92.

*E-mail: creignou@lim.univ-mrs.fr

†E-mail: daude@gypis.univ-mrs.fr

‡E-mail: Olivier.Dubois@lip6.fr

1 Introduction

For k -CNF formulas many experiments have shown a very swift transition between satisfiability and unsatisfiability as the ratio L/n of the number of clauses over the number of variables is varied. That is, there exists c_k , a critical value of L/n , such that if $L/n < c_k$ then the formula is almost surely satisfiable and if $L/n > c_k$ then the formula is almost surely unsatisfiable. Most of the papers investigating this phase transition are directed towards obtaining approximate estimates of its location. For instance for 3-SAT, for an observed sharp threshold of about $c_3 = 4.25$ the best lower bound is 3.003 [7] and tight upper bounds have been successively obtained, 4.601 [10], 4.596 [8] and finally 4.506 [5]. In 1999, Friedgut proposed a new and fruitful approach. In a remarkable paper [6], with an appendix by Bourgain, he developed a general sharp threshold criterion for monotone subsets of the hypercube. In using this criterion he proved that k -SAT [6] and (in collaboration with Achlioptas) k -col [1] exhibit a sharp threshold. In [4] Creignou and Daudé applied this criterion to the k -XOR-SAT problem (in which the usual “or” is replaced by the “exclusive or”). Thus, they proved the existence of a sharp threshold phenomenon for k -XOR-SAT, $k \geq 3$, without specifying its location.

The aim of this paper is to prove that the phase transition for k -XOR-SAT occurs as the number of clauses is proportional to the number of variables and more precisely to provide approximate estimates of its location. In Section 4 we give a lower bound and in Section 5 an upper bound. Our results rely first on the sharpness of the threshold established in [4] (see Section 2) and second on the handy translation of our problem in terms of random matrices (see Section 3). These theoretical results are supplemented with experiments in Section 6.

2 Notation and definitions

Throughout the paper k will denote an integer equal to or greater than 3. A k -XOR-clause (or shortly a k -equation), C , is a linear equation over the finite field $GF(2)$ using exactly k variables, $C = ((x_1 \oplus \dots \oplus x_k) = \varepsilon)$ where $\varepsilon = 0$ or 1. A k -XOR-formula (or shortly a k -system) is a conjunction of distinct k -XOR-clauses. A *truth assignment* I is a mapping that assigns 0 or 1 to each variable in its domain, it satisfies an XOR-clause $C = ((x_1 \oplus \dots \oplus x_k) = \varepsilon)$ iff

$I(C) := \sum_{i=1}^p I(x_i) \bmod 2 = \varepsilon$ and it satisfies a formula F iff it satisfies every clause in F .

We will denote by k -XOR-SAT (or shortly SAT) the property for a k -XOR-formula of being satisfiable (or equivalently the property for a k -system of being consistent) and by UNSAT the property of being unsatisfiable. The property UNSAT is monotone increasing.

Throughout the paper we reserve n for the number of variables ($\{x_1, \dots, x_n\}$ denotes the set of variables). There are $N_k = 2^{\binom{n}{k}}$ different k -XOR-clauses over n variables. We consider the random formula obtained by choosing uniformly,

independently and with replacement L clauses from the N_k possible k -clauses. This defines a probability space of k -XOR-formulas consisting of all ordered sets of L clauses, not necessarily distinct, with k distinct variables, each over a set of Boolean variables, each set of clauses having the same probability. This probability space is denoted by $\Omega(n, L, k)$, the associated probability is the uniform law:

$$\forall s \in \Omega(n, L, k) \quad P_{n,L}(s) = \left(2 \binom{n}{k} \right)^{-L}$$

We are interested in estimating the probability that a formula drawn at random from $\Omega(n, L, k)$ is satisfiable, that is in estimating $P_{n,L}(k\text{-XOR-SAT})$.

Our model for producing a random formula relates to the model in which each of the possible clauses is chosen independently with probability p in the same way $G(n, L)$ relates to $G(n, p)$ in random graph theory. In most investigations on properties of random subgraphs of the complete graph on n vertices, these two models are practically interchangeable, provided the number, L , of edges be close to $p \binom{n}{2}$ (see [3]). In [4] the first two authors proved that when each clauses is chosen with probability p then k -XOR-SAT exhibits a sharp threshold for $k \geq 3$. They also noted that the transition occurs when $pN_k \leq n$. Thus in the context of random formula these results are still valid in our model (provided $p \sim L/N_k$), they can be expressed as:

Theorem 2.1 [4]

For every $k \geq 3$ there exists a function $c_k(n) \leq 1$ such that: for every $\varepsilon > 0$

$$\lim_{n \rightarrow +\infty} P_{n, (c_k(n) - \varepsilon)n}(k\text{-XOR-SAT}) = 1$$

$$\lim_{n \rightarrow +\infty} P_{n, (c_k(n) + \varepsilon)n}(k\text{-XOR-SAT}) = 0$$

We are going to prove that this sharp threshold behavior occurs as the number of clauses is proportional to the number of variables of formulas and more precisely we are going to provide lower and upper bounds for $c_k(n)$. Since we know from Theorem 2.1 that $c_k(n) \leq 1$, we can and we will suppose in the sequel that $L \leq n$.

As mentioned in [6], though there is a swift transition of probability of satisfiability it is still feasible that the critical value $c_k(n)$ does not converge to any given value. However, one can define $\beta_k = \limsup_{n \rightarrow +\infty} c_k(n)$ and $\alpha_k = \liminf_{n \rightarrow +\infty} c_k(n)$.

Thus, β_k (respectively, α_k) is the least (greatest) real number such that if $c > \beta_k$ ($c < \alpha_k$) then the probability of a k -XOR-formula with n variables and cn clauses being satisfiable converges to 0 (to 1) as n tends to infinity; hence $\alpha_k \leq \beta_k$ and experiments suggest strongly (see Section 6) that equality holds.

In a first step (Section 3) we will reduce our problem to the study of the rank of random sparse matrices over $GF(2)$. Then, we will give a lower bound (Section 4) and an upper bound (Section 5) for the threshold.

At last, $H(x)$ will denote the well-known entropy function,

$$H(x) = x \ln(x) + (1 - x) \ln(1 - x).$$

3 From random formulas to random matrices

The aim of this section is to express $P_{n,L}(k\text{-XOR-SAT})$ in terms of random matrices. We will first use a technique related to the harmonic mean formula and publicized in [9] and [2]. This technique provides a simple expression for the number of satisfiable systems and can be described by a bipartite graph G formed by $\{0,1\}^n$ (the set of all assignments $I : \{x_1, \dots, x_n\} \longrightarrow \{0,1\}$) as the first part of vertices and $\Omega(n, L, k)$ (the set of all systems s) as the second part. There is an edge $\{I, s\}$ in G if the assignment I satisfies the system s . The degree of every assignment I is equal to $\binom{n}{k}^L$ and for every system s the degree $d(s)$ of s is the number of distinct assignments satisfying s . The number of satisfiable k -systems is thus given by

$$|k\text{-XOR-SAT}| = \sum_{I \in \{0,1\}^n} \sum_{s, I(s)=1} \frac{1}{d(s)}.$$

As there is a one-to-one correspondence from the set of linear systems satisfied by some fixed assignment I_0 onto the set $\mathcal{H}$ of homogeneous linear systems, we get:

$$|k\text{-XOR-SAT}| = 2^n \sum_{s \in \mathcal{H}} \frac{1}{d(s)}.$$

From this simple expression we get the following theorem.

Theorem 3.1 *Let $Y(A)$ denote the number of vectors in the kernel of the transpose of A , then providing A be chosen uniformly in the set $M_{L,n,k}$ of Boolean $L \times n$ matrices with exactly k units in each row:*

$$P_{n,L}(k\text{-XOR-SAT}) = \sum_{r=0}^L 2^{r-L} P_{(r)} = E_{L,k}(1/Y)$$

where $P_{(r)}$ is the probability that a matrix from $M_{L,n,k}$ is of rank r , and $E_{L,k}$ denotes the expectation.

Proof: From our preliminary work we have:

$$|k\text{-XOR-SAT}| = 2^n \sum_{A \in M_{L,n,k}} \frac{1}{2^{n-\text{rank}(A)}} = \sum_{r=0}^L \sum_{\substack{A \in M_{L,n,k} \\ \text{rank}(A)=r}} 2^r.$$

But $\#\{A \in M_{L,n,k} / \text{rank}(A) = r\} = |M_{L,n,k}| P_{(r)}$ and

$P_{n,L}(k\text{-XOR-SAT}) = |k\text{-XOR-SAT}| / \left(2 \binom{n}{k}\right)^L$, as $|M_{L,n,k}| = \binom{n}{k}^L$ we get:

$$P_{n,L}(k\text{-XOR-SAT}) = \sum_{r=0}^L 2^{r-L} P_{(r)}.$$

The second equality is justified by the following fact:

$$(\text{rank}(A) = r) \iff Y = 2^{L-r}.$$

■

We will see that this result leads to two key facts, Proposition 4.1 and Proposition 5.1, which enable us to give tight bounds for the threshold.

4 A lower bound for the threshold

By Theorem 3.1 and Jensen's inequality we have:

$$P_{n,L}(k\text{-XOR-SAT}) = E_{L,k}(1/Y) \geq \frac{1}{E_{L,k}(Y)}. \quad (1)$$

Thus, we get the first key fact:

Proposition 4.1 *If $\lim_n E_{cn,k}(Y) = 1$ then $\alpha_k \geq c$.*

A first lower bound for the threshold of k -XOR-SAT can be derived from the following result

Theorem 4.2 *Let θ_k denote the minimum over $[0, 1/2]$ of the function*

$$f_k(x) = \frac{\ln 2 + H(x)}{\ln(1 + (1 - 2x)^k)}$$

then for any $c < \theta_k$, $\lim_n E_{cn,k}(Y) = 1$

From MAPLE's estimates for θ_k we get the following bounds:

$$\alpha_3 > \mathbf{0.88949}, \quad \alpha_4 > \mathbf{0.96714}, \quad \alpha_5 > \mathbf{0.98916}, \quad \alpha_6 > \mathbf{0.99622} \quad (2)$$

To prove Theorem 4.2 we first show that the behavior of $E_{L,k}(Y)$ is given by the one of the following quantity:

$$\omega_{n,m,k} = \sum_{s=0}^k (-1)^s \binom{m}{s} \binom{n-m}{k-s}.$$

$$\mathbf{Proposition 4.3} \quad E_{L,k}(Y) = 2^{-n} \sum_{m=0}^n \binom{n}{m} \left(1 + \frac{\omega_{n,m,k}}{\binom{n}{k}}\right)^L.$$

Proof: Let us consider $X(A) = |\ker A|$, X is the sum of indicator variables:

$$X = \sum_{\vec{u} \in \{0,1\}^n} X_{\vec{u}} \text{ where } X_{\vec{u}}(A) = 1 \text{ if and only if } \vec{u} \in \ker A.$$

By symmetry, $E(X) = \sum_{m=0}^n \binom{n}{m} E(X_{\vec{u}_m})$ where $\vec{u}_m$ is any vector of weight m .

For each $\vec{u}_m$, consider the subset $I_{\vec{u}_m}$ of $\{1, \dots, n\}$ formed by the indexes of the unit coordinates of $\vec{u}_m$. Then $\vec{u}_m$ is in the kernel of A if and only if each row of the submatrix of A formed by the m columns whose indexes are in $I_{\vec{u}_m}$ has an even number s of unit coefficients. Thus, we have

$$\sum_{s=0 \bmod 2} \binom{m}{s} \binom{n-m}{k-s}$$

possible rows for a matrix A in $M_{n,L,k}$ such that $X_{\vec{u}_m}(A) = 1$, and we deduce:

$$\binom{n}{k}^L E(X_{\vec{u}_m}) = \left(\sum_{s=0 \bmod 2} \binom{m}{s} \binom{n-m}{k-s} \right)^L = \left(\frac{\binom{n}{k} + \omega_{n,m}}{2} \right)^L,$$

therefore, $E(X) = 2^{-L} \sum_{m=0}^n \binom{n}{m} (1 + \frac{\omega_{n,m,k}}{\binom{n}{k}})^L$.

As $Y = 2^{L-n} X$, the conclusion follows. ■

This result shows that if we split $E_{L,k}(Y)$ in two then the proof of Theorem 4.2 follows from the two following claims:

Claim 4.4 *If $L \leq n$ then $\lim_n \Sigma_1(L, k) = 1$, where*

$$\Sigma_1(L, k) = 2^{-n} \sum_{|m - \frac{n}{2}| < n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + \frac{\omega_{n,m,k}}{\binom{n}{k}})^L.$$

Claim 4.5 *If $c < \theta_k$ then $\lim_n \Sigma_2(\theta n, k) = 0$, where*

$$\Sigma_2(L, k) = 2^{-n} \sum_{|m - \frac{n}{2}| \geq n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + \frac{\omega_{n,m,k}}{\binom{n}{k}})^L.$$

These claims rely on asymptotical properties of sums of binomial coefficients and of the quantity $\omega_{n,m,k}$. Observe that for each k , $\omega_{n,m,k}$ can be viewed as a polynomial over two variables n and m of total degree k , this is made precise by the following:

Proposition 4.6

$$\omega_{n,m,k} = \frac{(n-2m)^k}{k!} - \frac{n(n-2m)^{k-2}}{2(k-2)!} + P_k(n, m)$$

where P_k is a polynomial on two variables of total degree $k-2$. Thus, there exists absolute positive constants A_k and B_k such that:

$$\left| \frac{\omega_{n,m,k}}{\binom{n}{k}} - \left(1 - \frac{2m}{n}\right)^k \right| \leq \frac{A_k}{n} \left|1 - \frac{2m}{n}\right|^{k-2} + \frac{B_k}{n^2} \quad (3)$$

Proof: Let us show how we get the leading term. Observe that:

$$\omega_{n,m,k} = \sum_{s=0}^k (-1)^s \frac{m^s}{s!} \frac{(n-m)^{k-s}}{(k-s)!} + P(n, m),$$

where $P(n, m)$ is a polynomial of degree $k-1$. But,

$$\sum_{s=0}^k (-1)^s \frac{m^s}{s!} \frac{(n-m)^{k-s}}{(k-s)!} = \frac{1}{k!} \sum_{s=0}^k \binom{k}{s} (-m)^s (n-m)^{k-s} = \frac{1}{k!} (n-2m)^k.$$

The study of the second term is left to the reader. ■

From this result and from its following consequences, Lemma 4.7 and Lemma 4.8, it turns out that the asymptotical behavior of $\Sigma_1(L, k)$ and $\Sigma_2(L, k)$ is given by the one of $(1 + (1 - \frac{2m}{n})^k)^L$.

Lemma 4.7 *If $L \leq n$ and $|m - \frac{n}{2}| < n^{1-\frac{4}{3k}}$ then*

$$\left| \left(1 + \frac{\omega_{n,m,k}}{\binom{n}{k}}\right)^L - \left(1 + \left(1 - \frac{2m}{n}\right)^k\right)^L \right| = O(n^{-7/9}),$$

$$\left| \left(1 + \left(1 - \frac{2m}{n}\right)^k\right)^L - 1 \right| = O(n^{-1/3}).$$

Proof: From (3) and from the mean value theorem we have when $L \leq n$:

$$\left| \left(1 + \frac{\omega_{n,m,k}}{\binom{n}{k}}\right)^L - \left(1 + \left(1 - \frac{2m}{n}\right)^k\right)^L \right| \leq \left(A_k \left|1 - \frac{2m}{n}\right|^{k-2} + \frac{B_k}{n} \right) \left(1 + \gamma_{n,m,k}\right)^{L-1}$$

with $\gamma_{n,m,k}$ lying between $\frac{\omega_{n,m,k}}{\binom{n}{k}}$ and $(1 - \frac{2m}{n})^k$.

As $|m - \frac{n}{2}| < n^{1-\frac{4}{3k}}$ and $k \geq 3$ we get $|1 - \frac{2m}{n}|^{k-2} = O(n^{\frac{-4}{9}})$. Thus, (3) gives $\left| \frac{\omega_{n,m,k}}{\binom{n}{k}} - \left(1 - \frac{2m}{n}\right)^k \right| = O(n^{\frac{-13}{9}})$ which shows that $\frac{\omega_{n,m,k}}{\binom{n}{k}} = O(n^{\frac{-4}{3}})$. Since we also have $|1 - \frac{2m}{n}|^k = O(n^{\frac{-4}{3}})$, we can conclude that $\gamma_{n,m,k}$ are $O(n^{\frac{-4}{3}})$. This leads to $\left(1 + \gamma_{n,m,k}\right)^{L-1} = O(n^{\frac{-1}{3}})$ and the rest of the proof is now pure routine. ■

Lemma 4.8 *If $L \leq n$, and $m - \frac{n}{2} \leq -n^{1-\frac{4}{3k}}$ or k even and $m - \frac{n}{2} \geq n^{1-\frac{4}{3k}}$ then*

$$\left| \frac{\left(1 + \frac{\omega_{n,m,k}}{\binom{n}{k}}\right)^L}{\left(1 + \left(1 - \frac{2m}{n}\right)^k\right)^L} \right| = O(1).$$

Proof: From (3) and from the mean value theorem we get:

$$\left| \ln\left(1 + \frac{\omega_{n,m,k}}{\binom{n}{k}}\right) - \ln\left(1 + \left(1 - \frac{2m}{n}\right)^k\right) \right| \leq \left(\frac{A_k}{n} \left|1 - \frac{2m}{n}\right|^{k-2} + \frac{B_k}{n^2} \right) \left(\frac{1}{1 + \nu_{n,m,k}} \right)$$

with $\nu_{n,m,k}$ lying between $\frac{\omega_{n,m,k}}{\binom{n}{k}}$ and $(1 - \frac{2m}{n})^k$.

Under the lemma's assumptions we get $(1 - \frac{2m}{n})^{k-2} \geq 2^{k-2} n^{-\frac{4(k-2)}{3k}}$.

Thus, (3) gives: $\frac{\omega_{n,m,k}}{\binom{n}{k}} \geq 2^{k-2} n^{-\frac{4(k-2)}{3k}} \left[4n^{\frac{8}{3k}} - A_k n^{-1} \right] - B_k n^{-2}$

which shows that there exists C_k such that for sufficiently large n

$$\frac{\omega_{n,m,k}}{\binom{n}{k}} \geq C_k n^{-4/3} > 0. \text{ As } (1 - \frac{2m}{n})^k > 0 \text{ we get } \nu_{n,m,k} > 0 \text{ and with } L \leq n,$$

the first inequality leads to: $\left| \frac{(1 + \frac{\omega_{n,m,k}}{\binom{n}{k}})^L}{(1 + (1 - \frac{2m}{n})^k)^L} \right| \leq \exp\left(A_k + \frac{B_k}{n}\right).$ ■

As mentioned above, the proof of Claims 4.4 and 4.5 will follow from well-known results on the behavior of the binomial coefficients and on the distribution of the binomial law.

Proof of Claim 4.4:

As $2^{-n} \sum_{m=0}^n \binom{n}{m} = 1$, Lemma (4.7) implies that:

$$\left| \Sigma_1(L, k) - 2^{-n} \sum_{|m - \frac{n}{2}| < n^{1 - \frac{4}{3k}}} \binom{n}{m} \right| = O(n^{-1/3}).$$

But, $k \geq 3$ hence $1 - \frac{4}{3k} > \frac{1}{2}$ and De Moivre-Laplace's theorem asserts that:

$$\lim_{n \rightarrow +\infty} 2^{-n} \sum_{|m - \frac{n}{2}| < n^{1 - \frac{4}{3k}}} \binom{n}{m} = 1.$$

■

Proof of Claim 4.5:

On the one hand, when k is odd and $m - \frac{n}{2} \geq n^{1 - \frac{4}{3k}}$, $-1 \leq 1 + (1 - \frac{2m}{n})^k \leq 2^{-k} n^{-\frac{4}{3}}$. Similar computations as in the proof of Lemma 4.8 show that for sufficiently large n , $-1 \leq \frac{\omega_{n,m,k}}{\binom{n}{k}} < 0$, thus $0 \leq (1 + \frac{\omega_{n,m,k}}{\binom{n}{k}})^L \leq 1$.

From De Moivre-Laplace's theorem we know that for any $k \geq 3$:

$$\lim_{n \rightarrow +\infty} 2^{-n} \sum_{|m - \frac{n}{2}| \geq n^{1 - \frac{4}{3k}}} \binom{n}{m} = 0.$$

Therefore, when k is odd, following Lemma 4.8

$$\Sigma_2(L, k) = \sum_{m - \frac{n}{2} \leq -n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + (1 - \frac{2m}{n})^k)^L + o(1).$$

On the other hand, when k is even Lemma 4.8 shows that

$$\Sigma_2(L, k) = O\left(\sum_{|m - \frac{n}{2}| \geq n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + (1 - \frac{2m}{n})^k)^L\right).$$

By parity and symmetry of binomial coefficients

$$\sum_{m - \frac{n}{2} \geq n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + (1 - \frac{2m}{n})^k)^L = \sum_{m - \frac{n}{2} \leq -n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + (1 - \frac{2m}{n})^k)^L.$$

Therefore, when k is even

$$\Sigma_2(L, k) = O\left(\sum_{m - \frac{n}{2} \leq -n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + (1 - \frac{2m}{n})^k)^L\right).$$

Therefore, Claim 4.5 is proved as soon as we are able to prove that for any positive real $c < \theta_k$:

$$T_{c,n} = 2^{-n} \sum_{m - \frac{n}{2} \leq -n^{1 - \frac{4}{3k}}} \binom{n}{m} (1 + (1 - \frac{2m}{n})^k)^{cn} = o(1). \quad (4)$$

From Stirling's formula (see [3]), we have for any $m \geq 1$,

$$\binom{n}{m} \leq \frac{1}{\sqrt{2\pi}} \left(\frac{n}{m(n-m)}\right)^{1/2} \exp\left(-n H\left(\frac{m}{n}\right)\right) e^{1/12n};$$

thus,

$$\binom{n}{m} = O\left(\exp\left(-n H\left(\frac{m}{n}\right)\right)\right).$$

Therefore,

$$T_{c,n} = O\left(\sum_{m - \frac{n}{2} \leq -n^{1 - \frac{4}{3k}}} \exp\left(-n g_{c,k}\left(\frac{m}{n}\right)\right)\right) \quad (5)$$

where

$$g_{c,k}(x) = \ln(1 + (1 - 2x)^k)(f_k(x) - c).$$

Now let us first note that for any $c < \theta_k$, $(f_k(x) - c) \geq \theta_k - c > 0$. Hence, for any $\alpha > 0$ there exists $\delta > 0$ such that:

$$g_{c,k}(x) > \delta \text{ when } x \in [0, 1/2 - \alpha].$$

Then, the Taylor expansion gives for any $k \geq 3$: $g_{c,k}(1/2 - t) = 2t^2 + o(t^2)$. Thus, there exists an absolute positive constant C_k such that

$$g_{c,k}(x) \geq C_k n^{-8/3k} \text{ when } x \in [1/2 - \alpha, 1/2 - n^{-4/3k}].$$

This proves the existence of some $K > 0$ such that when $m - n/2 \leq -n^{1-4/3k}$,

$$\exp(-n g_{c,k}(\frac{m}{n})) \leq \exp(-Kn^{1/9}),$$

with (5), this establishes (4) and thus Claim 4.5. $\blacksquare$

5 An upper bound for the threshold

Let Q denote the property “ A is of maximal rank” then $Pr(Q) = P_{(L)}$.

Since $P_{(L)} + P_{(L-1)} \dots + P_{(0)} = 1$, we have $\sum_{r=0}^L 2^{r-L} P_{(r)} \leq P_{(L)} + \frac{1 - P_{(L)}}{2}$.

Therefore, from Theorem 3.1 we get:

$$P_{(L)} \leq P_{n,L}(k\text{-XOR-SAT}) \leq \frac{1 + P_{(L)}}{2}.$$

Since we know that $k\text{-XOR-SAT}$ exhibits a sharp threshold (see Theorem 2.1) we get the second key fact:

Proposition 5.1 *The decreasing property Q exhibits a sharp threshold whose location coincides exactly with the one of $k\text{-XOR-SAT}$. Moreover if for some c there exist $\epsilon > 0$ such that for $L = cn$ and all (sufficiently large) n*

$$P_{(L)} < 1 - \epsilon \text{ then } \beta_k < c.$$

In other words, c is an upper bound for the threshold of $k\text{-XOR-SAT}$ as soon as we are able to prove that a matrix from $M_{cn,n,k}$ is not of maximal rank with positive probability.

When $k = 3$, such a bound can be derived from the following result:

Theorem 5.2

$$P_{(L)} \leq 1 - \frac{\exp(-3c) + 3c \exp(-6c) - c \exp(-9c) - 1 + c}{c} + o(1).$$

Therefore, solving the equation

$$\exp(-3c) + 3c \exp(-6c) - c \exp(-9c) - 1 + c = 0,$$

in using MAPLE provides

$$\beta_3 < \mathbf{0.9278}. \tag{6}$$

Proof: The matrices we consider, in $M_{L,n,3}$, have three units by row. Let $A = (a_{i,j})_{\substack{i=1,\dots,L \\ j=1,\dots,n}}$ be such a matrix. For each row $(a_{i,1}, \dots, a_{i,n})$, $i = 1, \dots, L$, of A there are three distinct indices $1 \leq i_1 < i_2 < i_3 \leq n$ such that $a_{i,j} = 1$ if $j = i_1, i_2$ or i_3 , and 0 otherwise; let $I_i(A)$ denote the set $\{i_1, i_2, i_3\}$. So, to each row we can associate its 3-weight, $W_i(A)$ defined as the following multiset:

$$W_i(A) = \left\{ \sum_{k=1}^L a_{k,i_1}, \sum_{k=1}^L a_{k,i_2}, \sum_{k=1}^L a_{k,i_3} \right\}.$$

Now, let us introduce three random variables T , U and V . For any random matrix A , let $T(A)$ count the number of all-zero columns in A . Observe that these columns do not contribute to the rank of A . Let $U(A)$ be defined by

$$U(A) = \#\{i / W_i(A) = \{1, 1, 1\}\}.$$

Observe that each row contributing to $U(A)$ underlines three columns such that only one of them contributes to the rank of A .

Finally, let $V(A)$ be defined by

$$V(A) = \#\{i / W_i(A) = \{1, 1, \alpha\} \text{ with } \alpha \geq 2\}.$$

Observe that each row contributing to $V(A)$ underlines three columns such that two of them contribute to the rank of A .

Thus, we have

$$\text{Rank}(A) \leq \text{Min}(L, n - T(A) - 2U(A) - V(A)).$$

Hence,

$$\text{if } T + 2U + V > n - L \text{ then } A \text{ is not of maximal rank, } L.$$

Therefore,

$$P_{(L)} \leq \text{Pr}(T + 2U + V \leq n - L). \quad (7)$$

Observe that

$$E(T + 2U + V) = \sum_{i=1}^n \text{Pr}(T + 2U + V \geq i).$$

Thus, for any integer $e \leq n$,

$$E(T + 2U + V) \leq e + (n - e) \text{Pr}(T + 2U + V \geq e + 1).$$

Hence,

$$\text{Pr}(T + 2U + V \geq e + 1) \geq \frac{E(T + 2U + V) - e}{n - e}.$$

Thus (7) gives for $L = cn$

$$P_{(L)} \leq 1 - \frac{\frac{E(T+2U+V)}{n} - 1 + c}{c}.$$

Let us estimate $E(T + 2U + V) = E(T) + 2E(U) + E(V)$.
The random variables T and U are the sum of indicator variables:

$$T = \sum_{j=1}^n T_j,$$

where $T_j(A) = 1$ iff the j th column of A is all-zero;

$$U = \sum_{1 \leq i_1 < i_2 < i_3 \leq n} U_{i_1, i_2, i_3},$$

where $U_{i_1, i_2, i_3}(A) = 1$ iff there exists i such that $I_i(A) = \{i_1, i_2, i_3\}$, and $W_i(A) = \{1, 1, 1\}$.

Now, $Pr(T_j = 1) = \left(\frac{\binom{n-1}{3}}{\binom{n}{3}} \right)^L$, hence

$$E(T) \geq n \exp(-3c)(1 - o(1)) \quad (8)$$

and

$$Pr(U_{i_1, i_2, i_3}) = L \frac{\binom{n-3}{3}^{L-1}}{\binom{n}{3}^L}, \text{ therefore,}$$

$$E(U) \geq cn \exp(-9c)(1 - o(1)). \quad (9)$$

Finally, let us estimate $E(V)$. In the same way

$$E(V) = \sum_{1 \leq i_1 < i_2 \leq n} E(V_{i_1, i_2}),$$

where $V_{i_1, i_2}(A) = 1$ iff there exists i such that $\{i_1, i_2\} \subset I_i(A)$, $\sum_{k=1}^L a_{k, i_1} = 1$, $\sum_{k=1}^L a_{k, i_2} = 1$, and $W_i(A) = \{1, 1, \alpha\}$ with $\alpha \geq 2$.

Let us introduce the random variables $Z_{i_1, i_2}(A)$ such that $Z_{i_1, i_2}(A) = 1$ iff there exists i such that $\{i_1, i_2\} \subset I_i(A)$, $\sum_{k=1}^L a_{k, i_1} = 1$ and $\sum_{k=1}^L a_{k, i_2} = 1$.

$$\begin{aligned} Pr(Z_{i_1, i_2} = 1) &= Pr(Z_{i_1, i_2} = 1 \text{ and } V_{i_1, i_2} = 1) + Pr(Z_{i_1, i_2} = 1 \text{ and } V_{i_1, i_2} = 0) \\ &= Pr(V_{i_1, i_2} = 1) + \sum_{i_3 \notin \{i_1, i_2\}} Pr(U_{i_1, i_2, i_3} = 1) \\ &= E(V_{i_1, i_2}) + (n-2)Pr(U_{i_1, i_2, i_3} = 1). \end{aligned}$$

But,

$$E(Z_{i_1, i_2}) = L(n-2) \frac{\binom{n-2}{3}^{L-1}}{\binom{n}{3}^L}.$$

Therefore,

$$E(V_{i_1, i_2}) = L(n-2) \frac{\binom{n-2}{3}^{L-1} - \binom{n-3}{3}^{L-1}}{\binom{n}{3}^L}.$$

And finally,

$$E(V) \geq (3cn \exp(-6c) - 3cn \exp(-9c))(1 - o(1)). \quad (10)$$

Finally, Theorem 5.2 follows from (8), (9) and (10). ■

The proof of Theorem 5.2 relies on a construction which can be carried out for all integer $k \geq 3$. For more readability we have restricted our attention to the generic case $k = 3$, however with similar considerations one can obtain the following general result :

Theorem 5.3 *for $k \geq 4$:*

$$P_{(L)} \leq 1 - \frac{\exp(-ck) + cu_k(c) + v_k(c) - 1 + c}{c} + o(1),$$

where

$$u_k(c) = \sum_{j=1}^{k-3} \binom{k}{j} \exp(-ck(k-j))((k-j+1) - (k-j-2)(k-j)),$$

$$v_k(c) = 3 \binom{k}{2} \exp(-2ck) + (-k^2 + 3k - 1) \exp(-ck^2).$$

Therefore, solving the equation

$$\exp(-ck) + cu_k(c) + v_k(c) - 1 + c = 0,$$

in using MAPLE provides

$$\beta_4 < \mathbf{0.9721}, \quad \beta_5 < \mathbf{0.9914}, \quad \beta_6 < \mathbf{0.9971} \quad (11)$$

6 Experiments

In this section, focussing on the generic case 3-XOR-SAT, we supplement the preceding rigorous results with experimental results. In the previous sections, we have determined an approximate scaling of a window in which a phase transition of the satisfiability must be observed, namely for a ratio number L of clauses to number n of variables below 0.8894, the probability of satisfiability tends to 1 as L and n tend to infinity and for a ratio above 0.9278, that probability tends to 0. In order to illustrate the phase transition and to estimate empirically the location of the critical value c_3 of the ratio for which the transition occurs, we have made experiments consisting in generating at random (in




Figure 1:

drawing uniformly and independently) 3-XOR-SAT formulas over 100, 200, 300 and 400 variables with a ratio varying from 0.70 to 1.14 in steps of 0.1. For each of these values of ratio, a sample of 1000 formulas has been solved with a computer program. The proportion of satisfiable formulas for each considered value of ratio has been plotted on the above Figure 1. It can be seen that the four smooth lines connecting the consecutive points corresponding to 100, 200, 300 and 400 variables, straighten as the number of variables increases showing thus strong empirical evidence of the sharp phase transition proved in [4]. The crossing of these lines suggest that the critical value c_3 of the transition is a little lower than 0.92 for 3-XOR-SAT.

7 Conclusion

The k -XOR-SAT problem is polynomial time solvable. Compared to the studies carried out on the phase transition of the SAT problem, this gives hope to get here an easier study. In a first step we have made precise the link between the k -XOR-SAT's phase transition and the rank of sparse random Boolean matrices. This last problem has been extensively studied by Russian mathematicians (see [11] and [12]), but reveals hard combinatorial and probabilistic problems. Our approach, which consists in using the sharpness of the threshold in order to specify its location, enables us to get good and interesting bounds for the critical ratio of k -XOR-SAT's phase transition. Indeed, experiments show the tightness of the upper bound. Moreover let us emphasize that Theorem 4.2 provides a new proof of Kolchin and Khokhlov's results [13], as well as a simpler expression for the corresponding critical value.

In conclusion, our work illustrates the importance of directing a lot of work towards obtaining general conditions for sharpness of a phase transition.

References

- [1] D. Achlioptas and E. Friedgut. A sharp threshold for k -colorability. *Random structures and algorithms*, 14(1):63–70, 1999.
- [2] D.J. Aldous. The harmonic mean formula for probabilities of unions: applications to sparse random graphs. *Discrete mathematics*, 76:167–176, 1989.
- [3] B. Bollobás. *Random graphs*. Academic Press, 1985.
- [4] N. Creignou and H. Daudé. Coarse and sharp thresholds for random k -XOR-CNF satisfiability. Submitted for publication to *Random Structures and Algorithms*, 2000.
- [5] O. Dubois, Y. Boufkhad and J. Mandler. Typical random 3-SAT formulae and the satisfiability threshold. *Proceedings of the 11th ACM-SIAM Symposium on Discrete Algorithms*, SODA’2000, pages 124–126, 2000.
- [6] E. Friedgut and an appendix by J. Bourgain. Sharp thresholds of graph properties, and the k -sat problem. *Journal of the A.M.S.*, 12(4):1017–1054, 1999.
- [7] A. Frieze and S. Suen. Analysis of two simple heuristics on a random instance of k -SAT. *Journal of Algorithms*, 20(2): 312–355, 1996.
- [8] S. Janson, Y.C. Stamatiou and M. Vamvakari. Bounding the unsatisfiability threshold for random 3-SAT. *Random Structures and Algorithms*, 17(2):103–116, 2000.
- [9] A. Kamatah, R. Motwani, K. Palem and P. Spirakis. Tail bounds for occupancy and the satisfiability threshold conjecture. In *Proceedings of the 35th Annual Symposium on Foundations of Computer Science*, Santa Fe, pages 592–603, 1994.
- [10] L.M. Kiroukis, E. Kranakis, D. Krizanc and Y.C. Stamatiou. Approximating the unsatisfiability threshold for random formulas. *Random Structures and Algorithms*, 12(3):253–269, 1998.
- [11] V.F. Kolchin. Random graphs and systems of linear equations in finite fields. *Random Structures and Algorithms*, 5(5):425–436, 1995.
- [12] V.F. Kolchin. *Random graphs*. Cambridge University Press, 1999.
- [13] V.F. Kolchin and V.I. Khokhlov. A threshold effect for systems of random equations of a special form. *Discrete Mathematics and Applications*, 2(5):563–570, 1992.